

Construction of a Non-2-colorable k -uniform Hypergraph with Few Edges

Heidi Gebauer *

November 5, 2009

Abstract

We show how to construct a non-2-colorable k -uniform hypergraph with $(2^{1+o(1)})^k$ edges. By the duality of hypergraphs and monotone CNF-formulas this gives an unsatisfiable monotone k -CNF with $(2^{1+o(1)})^k$ clauses.

1 Introduction

We will show the following.

Theorem 1.1. *For every $l \leq k$ we can construct a non-2-colorable k -uniform hypergraph with $m(k, l) = \binom{2^l - 1}{l} \cdot \left(\frac{2^l k}{l}\right)^l \cdot \left(\frac{2^l}{l} k\right)$ edges.*

The next proposition bounds $m(k, l)$

Proposition 1.2. *We have $m(k, l) \leq 2^{2l+l^2} \cdot k^l \cdot 2^k e^{\frac{k}{l}}$. In particular, $m(k, \log k) \leq (2^{1+o(1)})^k$.*

Hence we obtain a non-2-colorable hypergraph with few edges.

Corollary 1.3. *We can construct a non-2-colorable hypergraph with $(2^{1+o(1)})^k$ edges.*

Non-2-colorable hypergraphs connect to unsatisfiable CNF formulas: For a k -uniform hypergraph H let H' denote the k -CNF obtained by adding for every edge $e = (x_1, x_2, \dots, x_k)$ the clauses $C_e := (x_1 \vee x_2 \vee \dots \vee x_k)$ and $C'_e := (\bar{x}_1 \vee \bar{x}_2 \vee \dots \vee \bar{x}_k)$. Now H' is monotone, i.e., every clause either contains only non-negated literals or only negated literals. Moreover, every 2-coloring c of H yields a satisfying assignment α of H' (indeed, just set $\alpha(x_i) := 1$ if and only if x_i is colored blue under c) and vice versa. So Corollary 1.3 yields the following.

Corollary 1.4. *We can construct an unsatisfiable monotone k -CNF with $(2^{1+o(1)})^k$ clauses.*

*Institute of Theoretical Computer Science, ETH Zurich, CH-8092 Switzerland. Email: gebauerh@inf.ethz.ch.

2 Constructing a Non-2-Colorable Hypergraph with Few Edges

Throughout this section $\log$ stands for the binary logarithm. Moreover, a 2-coloring is an ordinary, not necessarily proper, 2-coloring.

Proof of Theorem 1.1: Let $k' = \frac{2^l}{l}k$. For every $i, i = 1, \dots, 2l - 1$, we let $A_i := a_{i,1}, a_{i,2}, \dots, a_{i,k'}$ be a sequence of length k' . Let c be a given 2-coloring. c has a *red majority* (*blue majority*) in the sequence A_i if under c at least $\frac{k}{2}$ elements of $\{a_{i,1}, a_{i,2}, \dots, a_{i,k'}\}$ are colored red (blue). Note that c has both a red majority and a blue majority in a sequence A_i if and only if there are equally many red and blue elements. We say that c has the *same majority* in the sequences $A_{i_1}, A_{i_2}, \dots, A_{i_j}$ if either c has a red majority in every sequence in $\{A_{i_1}, A_{i_2}, \dots, A_{i_j}\}$ or c has a blue majority in every sequence in $\{A_{i_1}, A_{i_2}, \dots, A_{i_j}\}$.

Proposition 2.1. *For every $\{X_1, \dots, X_l\} \subseteq \{A_1, A_2, \dots, A_{2l-1}\}$ we can construct a k -uniform hypergraph $G_{X_1, \dots, X_l}$ with at most $k^l \binom{k'}{l}$ clauses such that every 2-coloring c which has the same majority in $X_1, \dots, X_l$ yields a monochromatic edge in $G_{X_1, \dots, X_l}$.*

Proposition directly implies Theorem 1.1. Indeed, let G be the hypergraph consisting of the union of all edges in $G_{X_1, \dots, X_l}$ for every $\{X_1, \dots, X_l\} \subseteq \{A_1, A_2, \dots, A_{2l-1}\}$ and let c be a 2-coloring of the vertices of G . By the pigeon hole principle, for some $X_1, \dots, X_l \subseteq \{A_1, A_2, \dots, A_{2l-1}\}$, c has the same majority for $X_1, \dots, X_l$. But then c yields a monochromatic edge in $G_{X_1, \dots, X_l}$ and so c is not a proper 2-coloring of G . Since c was chosen arbitrarily G is not properly 2-colorable. Moreover, the number of edges of G is $\binom{2l-1}{l}$ times the number of edges in $G_{X_1, \dots, X_l}$, which gives the required number of edges in total. $\square$

Proof of Proposition 2.1: Let $X_j = x_{j,1}, x_{j,2}, \dots, x_{j,k'}$ for every $j, j = 1, \dots, l$. We will now shift sequences by a certain number of elements. For every $i \in \{0, \dots, k' - 1\}$ we let $X_j(i) = x_{j,1+i}, x_{j,2+i}, \dots, x_{j,k'}$, $x_{j,1}, \dots, x_{j,i}$. For every $i_1, i_2, \dots, i_l \in \{0, \dots, k' - 1\}$ and for every $S \subseteq \{1, 2, \dots, k'\}$ with $|S| = \frac{k}{l}$ we let $e_{i_1, i_2, \dots, i_l}(S)$ denote the set of elements which are of the form $x_{j,r+i_j}$ with $r \in S$. For every $i_1, i_2, \dots, i_l \in \{0, \dots, k' - 1\}$ we consider the hypergraph $G_{i_1, i_2, \dots, i_l} = \bigcup_{S \subseteq \{1, 2, \dots, k'\}: |S| = \frac{k}{l}} e_{i_1, i_2, \dots, i_l}(S)$. Let $G_{X_1, \dots, X_l}$ be the hypergraph consisting of the union of all edges in $G_{i_1, i_2, \dots, i_l}$ for every $i_1, i_2, \dots, i_l \in \{0, \dots, k' - 1\}$. Note that $G_{X_1, \dots, X_l}$ has $k^l \cdot \binom{k'}{l}$ edges, as claimed. It remains to show that every 2-coloring c which has the same majority in $X_1, \dots, X_l$ yields a monochromatic edge.

Proposition 2.2. *Let $s \in \{\text{red}, \text{blue}\}$ and let c be a 2-coloring which has an s -majority in X_i for every $i, i = 1, \dots, l$. Then there are $i_1, i_2, \dots, i_l$ such that for $\frac{k}{l}$ distinct r , $x_{1,r+i_1}, x_{2,r+i_2}, \dots, x_{l,r+i_l}$ all have color s under c .*

Proof: Choose $i_1, i_2, \dots, i_l$ uniformly at random from $\{0, 1, \dots, k' - 1\}$. For every r we let Y_r be the indicator variable for the event that $x_{1,r+i_1}, x_{2,r+i_2}, \dots, x_{l,r+i_l}$ all have color s under c . We

have $\Pr(Y_r = 1) \geq (\frac{1}{2})^l$. So the expected value $E[\sum_{i=1}^{k'} Y_i]$ is at least $k'(\frac{1}{2})^l = \frac{k}{l}$. Hence for some $i_1, i_2, \dots, i_l \in \{0, 1, \dots, k' - 1\}$, there are $\frac{k}{l}$ distinct r where $x_{1,r+i_1}, x_{2,r+i_2}, \dots, x_{l,r+i_l}$ all have color s under c . $\square$

Let $r_1, r_2, \dots, r_{\frac{k}{l}}$ be the distinct values for r described in Proposition 2.2. Let $S = \{r_1, r_2, \dots, r_{\frac{k}{l}}\}$. Then $e_{i_1, i_2, \dots, i_l}(S)$ is monochromatic under c . $\square$

Proof of Proposition 1.2: We use the following well-known fact. For every $r \leq n$,

$$\binom{n}{r} \leq \left(\frac{en}{r}\right)^r \quad (1)$$

By (1), $\binom{\frac{2^l}{l}k}{\frac{k}{l}} \leq (e2^l)^{k/l} = 2^k e^{k/l}$. Hence $m(k, l) \leq 2^{2l} \cdot 2^{l^2} \cdot k^l \cdot 2^k e^{\frac{k}{l}}$. Since $k^{\log k} = 2^{\log^2 k}$ we get $m(k, \log k) \leq (2^{1+o(1)})^k$. $\square$